

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C12Q 1/04, 1/70, 1/66	A1	(11) International Publication Number: WO 94/06931 (43) International Publication Date: 31 March 1994 (31.03.94)
(21) International Application Number: PCT/GB93/01989 (22) International Filing Date: 22 September 1993 (22.09.93) (30) Priority data: 9220027.8 22 September 1992 (22.09.92) GB (71) Applicant (for all designated States except US): THE MINISTER OF AGRICULTURE FISHERIES AND FOOD IN HER BRITANNIC MAJESTY'S GOVERNMENT OF THE UNITED KINGDOM OF GREAT BRITAIN AND NORTHERN IRELAND [GB/GB]; Whitehall Place, London SW1A 2HH (GB). (72) Inventor; and (75) Inventor/Applicant (for US only) : SANDERS, Michael, Frederick [GB/GB]; 9 Essex Road, Basingstoke, Hampshire RG21 1TA (GB).		(74) Agent: LOCKWOOD, Peter, Brian; MOD(PE), DIPR, 1PR 1b, Room 2002, Empress State Building, Lillie Road, London SW6 1TR (GB). (81) Designated States: AT, AU, BB, BG, BR, CA, CH, CZ, DE, DK, ES, FI, GB, HU, JP, KR, LK, LU, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: METHOD AND KITS FOR DETECTION OF BACTERIA (57) Abstract A method for detection, identification and/or quantification of target organisms of specific bacterial genus, species or serotype, based upon the occurrence of release of cell contents, particularly nucleotides e.g. ATP, on lysis of bacterial cell walls on incubation with bacteriophages (phages) specific for them. When new phage particles are released at the end of the phage replication cycle nucleotide levels are measured and compared with controls. The method provides for the detection of specific bacteria in which insertion of the lux gene into the phage genome is not required yet it is faster and more sensitive than known non-modified phage utilising techniques. The method is only limited by the availability of phage types suitable for selective attack of the target bacteria to be detected and can detect a single <i>Salmonella</i> in a sample of milk in under 12 hours.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TG	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

METHOD AND KITS FOR DETECTION OF BACTERIA.

The present invention relates to a method of detection, identification and/or quantification of bacteria and to test kits for carrying it out. Particularly the method enables detection of organisms of specific bacterial genus, species or serotype, in isolated form or as contaminants in environmental or forensic samples, or in foodstuffs.

There are many requirements for methods of screening for specific bacteria, particularly those present in low numbers and in specific environments; for example, human bacterial pathogens in contaminated foods. Public health and quality control bodies demand rapid bacterial detection methods which have suitable levels of specificity and sensitivity, but few satisfactory methods exist.

It is known to detect specific bacteria by use of genetically engineered bioluminescent bacteriophages, virus particles which have had the 'lux' gene inserted into their genome, as described by Ulitzer and Kuhn, in Scholmerich et al (Eds.) 'Bioluminescence and Chemiluminescence-new perspectives' pages 463-472: pub. (1987) by John Wiley and Sons. The 'lux' gene is that encoding for bacterial luciferase and the technique is based upon the fact that upon infection of a target bacterium, bacteriophage genes and the lux gene are injected into it and are subsequently expressed. The presence of the target bacterium is indicated by emission of light from the sample which can easily be measured. Most bacteria are susceptible to attack by bacteriophages (commonly called phages), many of which lyse or disrupt their host at the end of their replication process, and these interactions show varying degrees of host/phage specificity.

Schutzbank et al have shown the potential of this technique but note problems with cross reactivity between construct phages and other non-target bacterial types. While such problems may be overcome by engineering more specific phages, see for example US4348478, this

entails provision of phages for each type of target bacteria for which a need to test exists. Such recombinants may not readily be constructed for a variety of reasons, not least of which being the need to avoid disruption of the function of the phage itself.

Initially only Salmonella and E. coli phages were successfully modified. This is because a great deal of work had been previously done on the molecular biology of these organisms and it was known that the lux genes could fit into the phage head without causing loss of the phage's ability to be infective. This is not the case with other food pathogens and a massive amount of work will need to be done before a comprehensive range of modified phages can be constructed.

A further method utilising the specificity of phages for their host bacteria is disclosed by Hirsh and Martin (1984) Journal of Food Protection, Vol 47, No 5, pp388-390. This method relies on the phage induced destruction of the bacteria grown on enrichment broth to produce lacunae in the bacterial lawns, wherein lacunae are indicative of the presence of the target bacterium; in this case a Salmonella. However the method is lengthy to perform, requiring about 24 hours for a positive result, and is only sensitive to 5 cells/ml or more.

The present inventor has now provided a method based upon the occurrence of release of cell contents on damage, eg. lysis, of cell walls of bacteria when new phage particles are released at the end of the phage replication cycle. This method provides for the detection of specific bacteria which does not require insertion of the lux gene into the phage genome, whereby specificity is only limited by the availability of phage types suitable for selective attack of the target bacteria to be detected, yet its performance takes far less time than the existing unmodified phage techniques. Thus the method of the present invention has more readily realised potential for the specific and rapid detection of almost any bacteria in any environment; (foodstuffs, drinking water, pharmaceutical products and diseased

tissues in humans, animals and plants etc.) provided that a phage with the necessary specificity can be found. Use of the method of the present invention has been shown to be capable of detection of a single Salmonella in a 1ml sample of milk in under 12 hours.

Thus the present invention provides a method for the detection, identification and/or quantification of target organisms of specific bacterial genus, species or serotype present in a material comprising incubating the material or bacteria derived therefrom with bacteriophage selected for the ability to specifically infect target organisms and release cellular components from them, measuring the amount of one or more particular components released from any bacteria present during the incubation, and relating this to the presence, identity and/or amount of target organisms. Preferably incubation of material with bacteriophage is carried out in bacterial support medium.

Preferred cellular component comprises nucleotides. Theoretically it is possible to measure any of the nucleotides that are released by the cell lysis caused by the release of new phage particles, for example NAD, NADP, NADH, NADPH, ATP or ADP, cAMP or cGMP, with sensitivity provided by use of one or more of the many enzyme based assay systems, eg. 'cascade' systems, that are available in the art. For example, GB 2213261 discloses a method which may be used for assaying reduced pyridine nucleotides, eg NADH or NADPH, based upon a salicylate monooxygenase system, while other enzyme systems such as the alkaline phosphatase/NAD/NADP system as disclosed in GB 2240845. Suitable assays for ADP, cAMP, cGMP etc will occur to those skilled in the art.

However, particularly preferred is the measurement of adenosine triphosphate (ATP), that being readily measurable by assay with a variety of enzyme/enzyme substrate combinations by virtue of its being a cofactor in numerous substrate conversions, and being released in relatively large quantities as compared with other bacterial nucleotides. For the rapid and efficient determination of levels of

released ATP it is especially preferred to utilise enzyme reactions which result in production of luminescence, most conveniently using luciferase. ATP release is quantifiable with commercially available reagents using bioluminescence wherein it is used to drive oxidation of luciferin under catalysis by luciferase resulting in the emission of light. The quantum efficiency of this reaction is extremely high and the presence and amount of light produced gives a measure of ATP released and thus of the presence and numbers of target organisms.

For identification or quantification of specific bacteria occurring in material in relatively high concentrations, eg. in cultures of isolated bacteria, it is possible merely to incubate the specific phage with a part of that culture or a subculture in the presence of, or with subsequent addition of, the component assay reagents and thus to measure the amount of component released by performing the assay.

For identification, detection and/or quantification of specific bacteria at lower concentrations, for example as contaminants in or on water or foodstuff materials, it is necessary to first perform an enrichment of bacteria derived from the material under test, eg. for a few hours, to allow the target bacterium to multiply to a level where its released components, eg. nucleotides, will be detectable above background levels. Enrichment is preferably carried out by culturing a selective medium, favouring growth of target bacteria, that has been inoculated with the material or a culture derived therefrom, eg. derived from a swab. Typically enrichment lasts less than a working day, eg. 1 to 10 hours; preferably 1 to 5 hours.

After enrichment, one or more phage types with known target host specificities are mixed with the culture and the mixture is incubated. During the incubation period infection of specific target bacteria with phage occurs and the replication cycle begins. At the end of the replication cycle target bacteria present burst (times range from eg.

20 - 60 minutes depending on species), cellular component, eg. nucleotide, is released and detected. Control samples either containing no bacteria, non-target bacteria or bacteria without phage show no substantial increase in levels above background levels.

Furthermore, dependent upon the biology of the phage/host pair selected, the method has the potential for great specificity and still further sensitivity and rapidity. Utilising the aforesaid ATP assay method the inventor has readily developed systems for the detection of several human pathogens in milk including Staphylococcus aureus, Listeria monocytogenes and Salmonella, and others for foodstuffs such as lettuce whereon pathogens such as Listeria are known to be a potential danger. Salmonellae, Listeria, Staphylococci, E. coli and pseudomonads have all proved to be readily detectable by the method. It will be seen by those skilled in the art however that there is no limit to the application of the present method other than the availability of the necessary specific phage or phages.

In a more preferred embodiment of the present invention it has been found to be advantageous to incubate a sample of material under investigation with a nonselective bacterial support medium, eg. such as peptone water or Brain Heart Infusion broth (B.H.I), in order to activate or 'resuscitate' substantially all of the bacteria present therein, before carrying out any enrichment incubation. Such incubation is conveniently, but not essentially, carried out for 1-5 hours; but as with the enrichment, shorter or longer periods may be appropriate depending on, inter alia, the freshness of the material.

In a preferred embodiment a sample of the resuscitated material is then transferred to a selective medium adapted to favour growth of the bacteria being investigated over interfering bacteria and incubated for a further period, eg. 1-5 hours. Conveniently an aliquot of the medium from the resuscitation stage is so transferred.

After the incubations are complete all or part of the cultured media

is mixed with the selected phage, eg. phage in the form of a suspension or freeze dried, and incubated for a set time and at a set temperature selected according to the biological characteristics of the phage/bacteria combination being used. The amount of component, eg. nucleotide, whose release is being measured is then measured using an appropriate assay, eg. enzyme/substrate system.

For assaying the release of cellular component completed resuscitation or selective enrichment step medium, or a medium with a sample of the material itself, is conveniently mixed with the selected phage in a vessel, eg for ATP in a luminometer tube, and incubated for a suitable time and suitable temperature, eg. 30 to 60 minutes at 20°C to 40°C, to allow cell lysis; these conditions depending on the phage/bacteria system. For measuring ATP, light producing assay reagents, eg. those of the luciferin/luciferase system, can be added before or after this incubation period or the lysis period; light produced over the period or on addition is detected in a luminometer and related to the amount of ATP released. In all cases controls are advantageously carried out for comparison of background component levels, eg. nucleotide levels, in samples incubated without the phage or incubated with phage and a known amount of target bacteria whereby production of calibration curves for assessing bacteria numbers is enabled. Such controls may include challenge with other phages of different specificity in order to more completely determine the characteristics of the organisms present in the material. Similarly several phage types may be used in the same incubation where the method is being used to screen for a number of types of bacteria for which no one common phage is specific.

The range of bacteriophages available and the bacteria for which they are specific will be realised to be vast by those skilled in the art. For example a list of phage types available from the American Type Culture Collection (ATCC) is published by them as 'Catalogue of Bacteria & Bacteriophages'. Other such depositories also publish equivalent data in their catalogues and this may be used to identify

possible phage 'reagents' for the present method. Phages may be used, inter alia, in aqueous suspension or in freeze dried form eg. on microtitre plate wells. In this manner plate luminometry can be used. Preferred phages are lytic and lead to lysis of the target organism.

In addition to deposited phages, a further source of phages is provided by isolating them from suitable environments; that is conveniently the environment where the target bacteria are themselves to be found. For example, it is possible to isolate phages specific to both Campylobacter spp. and Salmonella spp. from effluent from a poultry processing plant. Isolation techniques will be well known to those skilled in the art but are exemplified by Loessner and Busse (1990) Applied and Environmental Microbiology Vol 56, pp1912-1918, and Adams 'Bacteriophages' Pub Interscience Inc (1959) pp447-455.

The range of media available for selective promotion of growth of a particular bacterial type will also be known to those skilled in the art and these may function by positive action or by eg. inhibition of other organisms. Examples of such media are illustrated by reference to supplier's manuals, eg. such as those available from UNIPATH Limited, Wade Road, Basingstoke, HANTS, RG24 0PW, UK 'Selective Microbiology for Food and Dairy Laboratories', or eg. the OXOID manual. These publications list, for example, media capable of favouring growth of Campylobacter, Listeria and Yersinia. Similarly methods for isolation of food pathogens for preparation of test samples are well known. (UNIPATH and OXOID are Registered Trade Marks).

The present invention also provides test kits for carrying out the method of the present invention and these are characterised in so far as they comprise a bacteriophage selected for the ability to specifically infect bacterial target organisms, ie. a type or types for which detection, identification and/or quantification is desired, in combination with some or all of the reagents which are specifically associated with the aforesaid method of the invention.

Thus preferably test kits of the present invention comprise;
(a) a bacteriophage selected for its ability to specifically infect target bacterial organisms and release cellular components from them, preferably in a form substantially free of such target organisms, and
(b) the reagents necessary for carrying out assay of the amount of one or more cellular components released by the action of the bacteriophage on the bacterial organism.

Optionally the kits further comprise (c) a selective medium for promoting preferential growth of target bacterial organisms and/or (d) a non-selective bacterial growth medium.

Thus preferred test kits of the invention are those wherein the reagents necessary for carrying out assay are for assay of the amount of a nucleotide present in a sample, preferably comprising luciferin and luciferase. Test kits optimised for performance of high sensitivity assay of bacteria will include both non-selective and selective growth media, the non-selective growth medium being preferably that suitable for a microorganism resuscitation step.

It will be realised that, unlike the prior art method which monitors lacunae on solid media supported bacterial lawns, the present method is particularly adapted to test for bacteria derived from samples of material under investigation by culturing on liquid media, although solid media may of course be used. A particular kit for performance of the present method may be one having the bacteriophage component (a) together with one or more media (c) and/or (d) of liquid form or concentrated or dry components adapted for the preparation of such liquid medium. While many such combinations adapted for the performance of the method of the present invention will occur to those skilled in the art, the most complete form of the kit will comprise all the reagents (a) to (d); the nucleotide assay components preferably being optimised for measuring nucleotide levels in a sample of the bacterial growth medium. Where quantification is desired

samples of the bacteria might also be included, eg. as spores or in cells in a medium, for setting up control readings of known levels of bacteria for calibrating the nucleotide assay. Target and control non-target organisms may be used.

The method and kits of the present invention will now be exemplified by way of illustration only by reference to the following non-limiting examples. The vast variety of options available will be readily determinable by those skilled in the art on consideration of the general method described above and particularised below, and the available types of bacteriophages and nucleotide assays.

EXAMPLE 1. Protocol and test kit for detection of low levels of Salmonella in milk by phage-mediated ATP bioluminescence.

Strains used: the strain of bacterium to be detected was a Salmonella which had been isolated from poultry processing plant effluent water. Prior to investigations stock cultures of bacteria were maintained on Brain Heart Infusion (B.H.I.) agar (Unipath Ltd., Basingstoke, Hampshire) slants. The specific bacteriophage, Sall, was also isolated from this source.

The bacteriophages were stored as bacteria-free lysates in 1.5ml cryotubes above liquid nitrogen; the titre of bacteriophage lysate used was 1.2×10^9 plaque-forming units (p.f.u.) per millilitre.

Brain Heart Infusion broth (BHI) -Unipath Ltd. was used to grow the inoculum. Serial dilutions of broth culture were made using 0.1M potassium phosphate buffer (pH 7). Selective enrichment was shown to be achieved using Rappaport-Vassiliadis (R.V.) broth (Unipath Ltd.), and more preferably Selenite Cystine (S.C) broth (Difco Laboratories, Detroit, Michigan).

A suitable kit for detection of this Salmonella using the method of

the present invention was provided comprising (a) specific bacteriophage Sall, (b) luciferin/luciferase assay reagents (eg. as below) and/or (c) R.V. or S.C. broth and optionally included (d) B.H.I. Sall is optionally provided as infected bacteria which are completely lysed to liberate phage and destroy bacteria before use.

The spectrophotometer used was model CE 202 (Cecil Instruments, Cambridge, England) with 1 cm disposable cuvettes (Whatman International Ltd., Maidstone, England). Light measurements were made using a luminometer, model LB953 Autolumat (Berthold Instruments U.K. Ltd., St. Albans, Hertfordshire), and disposable polystyrene tubes (catalogue number 55.476; Sarstedt, Beaumont Leys, Leicester, U.K.). The adenosine 5'-triphosphate assay reagents were luciferin/luciferase and adenosine 5'-triphosphate assay mix dilution buffer and were obtained from Sigma Chemical Co Limited, Poole, Dorset. Luciferin /luciferase reagent was diluted 1:25 with the buffer when required.

Preparation of inoculum: A broth culture of Sall was set up in 10 ml B.H.I. broth and incubated at 37°C until visibly turbid. Absorbance at 650 nm was measured, and simultaneously a tenfold dilution series from 10^0 through to 10^{-10} was prepared in potassium phosphate buffer. From each dilution of culture five 1 ml samples were inoculated with fresh 9 ml volumes of B.H.I. broth. These tubes were then incubated overnight at 37°C, and observed for turbidity. From published tables ("Theory and Practice in Experimental Bacteriology", G.G. Meynell & Elinor Meynell, Cambridge University Press) it was possible to determine which culture dilution contained approximately 100 cells per millilitre and that dilution was used to prepare samples for testing.

i) Protocol: Milk test: sampling and enrichment procedure A 1ml sample of 100 ml fresh pasteurised milk infected with approximately 1 cell per millilitre of log-phase Sall at room temperature was transferred into 9 ml fresh B.H.I. broth and incubated for 2 hours at 37°C as a resuscitation/enrichment step. Either (a) 0.1 ml of this

medium was then placed in 10 ml Rappaport-Vassiliadis (RV) selective -enrichment broth and incubated at 41°C for 7 hours.

Alternatively, where it was suspected that the viability of bacteria in the sample is poor, a less potent selective medium was applied, as death of weakened bacterial cells was recorded in presence of R.V. medium in some cases. The 'milder' medium was selected was Selenite Cystine broth wherein (b) 1.0 ml of the completed resuscitation /enrichment step medium was placed in 9ml Selenite Cystine selective enrichment broth and incubated at 35°C overnight, with the 1 ml of the resultant culture used to inoculate 9ml fresh B.H.I. which was then incubated at 37°C for 2 hours as a recovery step.

In either case (a) or (b), 4.5 ml of the completed culture was mixed with 0.5 ml specific bacteriophage suspension Sal1. The remainder of the R.V., Selenite Cystine or B.H.I recovery culture was retained.

(ii) Preparation of luminometer and light measurement 50 microlitres B.H.I., held at 37°C, was added to all 80 polystyrene luminometer tubes in the luminometer, which itself had been pre-warmed to 37°C. 50 microlitres of the R.V. or B.H.I cultures, dependent upon the protocol selected, to which bacteriophages had been added was placed in each alternate tube in the luminometer, and 50 microlitres of the respective culture without bacteriophages was added to the remaining tubes as the negative control. Light measurement was then commenced, with 100 microlitres of luciferin/luciferase reagent injected into each tube and light output immediately monitored over periods of 60 seconds. Measurements were then plotted as peak light intensity against time for both positive and negative control tubes.

Using either protocol a peak of 25 000 counts per second (c.p.s.) was obtained after 60 minutes for positive tubes, with the negative controls reaching only 500 c.p.s.

EXAMPLE 2. Protocol and test kit for the detection of *Listeria monocytogenes* in lettuce by phage mediated ATP bioluminescence.

Strains used: The strain of *Listeria monocytogenes* used was ATCC 23074 with its specific bacteriophage ATCC 23074-B1. The bacteriophage was stored as a bacteria-free lysate in 1.5 ml cryotubes above liquid nitrogen. The titre of the phage preparation was approximately 2.11×10^{10} p.f.u. per millilitre.

Culture media: Stock cultures were maintained on Brain Heart Infusion (B.H.I.) agar slants. B.H.I. broth was used for growth of inocula. Serial dilutions were made using potassium phosphate buffer (0.1 M, pH7). The selective-enrichment medium was *Listeria* Enrichment Broth (L.E.B.). All culture media were obtained from Unipath Ltd., Basingstoke, Hampshire, U.K.

A suitable kit for the determination of *Listeria monocytogenes* by the method of the present invention was provided comprising (a) bacteriophage ATCC 23074-B1, (b) the luciferin/luciferase assay reagents and/or (c) L.E.B and optionally included B.H.I.

Apparatus: The spectrophotometer, cuvettes, luminometer and polystyrene luminometer tubes were as detailed in the salmonella protocol. A Colworth 400 stomacher was used (A.J. Seward, London SE1).

Reagents: Luminescence reagents were as described in the *Salmonella* protocol.

Preparation of inoculum: An inoculum of ATCC 23074 containing approximately 100 cells per millilitre was prepared in the same way as described in the *Salmonella* protocol.

Inoculation of lettuce for use as positive test: Fresh lettuce was chopped by hand into pieces approximately 2 centimetres in length.

100 mls of the buffer suspension of Listeria prepared in step (i) was added to 100 grams of chopped lettuce and mixed well by shaking for 1 minute. The lettuce was then drained and stored at 4°C.

i) Extraction and enrichment procedures for Listeria: The inoculated lettuce was placed in a large stomacher bag with 100 mls potassium phosphate buffer and mixed in the stomacher for 1 minute. 1 ml of the resultant suspension was then added to 9 ml B.H.I. broth and incubated at 30°C for 4 hours as a resuscitation/enrichment step. 1 ml of this culture was then added to 9 ml L.E.B. at 30°C and incubated overnight, this being the selective-enrichment step. 4.5 ml of this culture was then removed and 0.5 ml of specific bacteriophage ATCC 23074-B1 was added to that.

The remaining L.E.B. culture was kept as the negative control.

ii) Preparation of luminometer and light measurement: 50 microlitres warm (30°C) B.H.I. was added to 80 tubes in the luminometer, which had been pre-warmed to 30°C. The machine was set to run in kinetics mode at 30°C. To the first tube, and to each alternate tube after it, was added 50 microlitres of the negative control culture (without bacteriophages). To the remaining tubes was added 50 microlitres of the test culture (with bacteriophages). 100 microlitres luciferin /luciferase reagent was then injected into each tube in turn and the light output immediately measured over periods of 60 seconds.

As with Salmonella, the peak light intensity (counts per second) for the test and negative control tubes was plotted against time. Following this protocol, a peak of 10 000 c.p.s. was observed for the test culture after 48 minutes whilst the negative control remained constant with approximately 1000 c.p.s. The absence of light output levels above the control is interpreted as absence of target Listeria organisms and not absence of bacteria per se.

EXAMPLES 3 to 11.

Further examples of bacteria and particular phage that are capable of specifically releasing their cell contents, including ATP, are listed below with media and preferred incubation temperature suitable for incubating these together for the purpose of the present method.

EXAMPLE No	BACTERIA	PHAGE	MEDIUM	TEMP
3	Lactococcus lactis sp cremoris ATCC11603	ATCC 11603 -B1	M17 Broth Unipath	26°C
4	Enterococcus faecium ATCC19950	ATCC 19950 -B1	M17 Broth Unipath	37°C
5	Staphylococcus aureus subsp. aureus ATCC 11988	ATCC 11988 -B1	BHI (Oxoid) CM 225 Unipath	37°C
6	Bacillus cereus ATCC 12826	ATCC 12826 -B1	BHI (Oxoid) CM 225 Unipath	30°C
7	'' ATCC 27877	ATCC 27877 -B1	BHI (Oxoid) CM 225 Unipath	30°C
8	Enterobacter cloecae ATCC 23355	ATCC 23355 -B1	BHI (Oxoid) CM 225 Unipath	30°C

EXAMPLE No	BACTERIA	PHAGE	MEDIUM	TEMP
9	Staphylococcus aureus subsp. aureus ATCC 11987	ATCC 11987 -B1	BHI (Oxoid) CM 225 Unipath	37°C
10	Shigella dysenteriae (Shiga) Castellani & Chalmers ATCC 11456a	ATCC 11456a -B1	BHI (Oxoid) CM 225 Unipath	37°C
11	Shigella dysenteriae ATCC 23351	ATCC 23351 -B1	BHI (Oxoid) CM 225 Unipath	37°C

CLAIMS.

1. A method for the detection, identification and/or quantification of target organisms of specific bacterial genus, species or serotype present in a material comprising incubating the material or bacteria derived therefrom with bacteriophage selected for the ability to specifically infect target organisms and release cellular components from them, measuring the amount of one or more particular cellular components released from any bacteria present during the incubation, and relating this to the presence, identity and/or amount of the target organism.
2. A method as claimed in claim 1 wherein the incubation is carried out in a bacterial support medium.
3. A method as claimed in claim 1 or 2 wherein the incubation is carried out at between 20°C and 40°C.
4. A method as claimed in claim 1, 2 or 3 wherein the one or more cellular components comprise a nucleotide selected from NAD, NADP, NADH, NADPH, ATP, ADP, cAMP or cGMP.
5. A method as claimed in claim 3 or 4 wherein the nucleotide is measured using an enzyme based assay system.
6. A method as claimed in any one of the aforesaid claims wherein the component is ATP and the assay is one that utilises an enzyme resulting in the production of luminescence.
7. A method as claimed in claim 6 wherein the assay system is the luciferin/luciferase system and the amount of light given off is measured over the entire incubation or over the cell lysis period.
8. A method as claimed in any one of the preceding claims wherein the step of incubating the material or derived bacteria with the specific

bacteriophage is carried out in the presence of, or with subsequent addition of, the particular cellular component assay reagents.

9. A method as claimed in any one of claims 1 to 8 wherein the material or derived bacteria is/are incubated with a selective medium favouring growth of the target bacteria, or a non-selective medium capable of supporting bacterial growth, prior to the incubation with the bacteriophage.

10. A method as claimed in claim 9 wherein the incubation with selective medium is for 1 to 10 hours.

11. A method as claimed in any one of the preceding claims wherein the incubation with the bacteriophage is for between 20 and 60 minutes.

12. A method as claimed in any one of the preceding claims wherein prior to the incubation with selective medium or with the bacteriophage, the material or bacteria is/are incubated with a non-selective medium, and on completion of this incubation a sample of the medium is transferred to the selective medium or bacteriophage incubation step as source of material or bacteria to be tested.

13. A method as claimed in claim 12 wherein the incubation with non-selective medium is for between 1 and 5 hours.

14. A method as claimed in any one of the preceding claims wherein the bacteriophages are provided in the form of a suspension, as freeze dried phage, or as infected bacteria which are lysed before use.

15. A method as claimed in any one of the preceding claims wherein several bacteriophage types are used in the same incubation and the method is used to screen for a number of types of bacteria.

16. A test kit for the detection, identification and/or quantification of target organisms of specific bacterial genus, species or serotype comprising a bacteriophage selected for its ability to specifically infect target organisms and release cellular components from them, characterised in that the test kit further comprises some or all of the reagents which are specifically associated with the method of any one of the claims 1 to 15.
17. A test kit for the detection, identification and/or quantification of target organisms of specific bacterial genus, species or serotype comprising
- (a) a bacteriophage selected for its ability to specifically infect target bacterial organisms and release cellular components from them, and
 - (b) the reagents necessary for carrying out assay of the amount of a cellular component released by the action of the bacteriophage on the bacterial organism.
18. A test kit as claimed in claim 17 further comprising (c) a selective medium for promoting preferential growth of organisms of bacterial genus, species or serotype for which the kit is targeted and/or (d) a non-selective bacterial growth medium.
19. A test kit as claimed in claim 17 or 18 wherein the cellular component to be assayed by reagent (b) comprises a nucleotide.
20. A test kit as claimed in claim 19 wherein the reagents necessary for carrying out assay of the amount of a nucleotide present in a sample comprise luciferin and luciferase.
21. A test kit as claimed in any one of claims 17 to 20 comprising the medium (c) and/or (d) in liquid form or as concentrated or dry components required for the preparation of such liquid medium.

22. A test kit as claimed in any one of claims 16 to 21 further comprising target organisms and/or control non-target organisms.

23. A test kit as claimed in any one of claims 16 to 21 substantially as described in either of Example 1 or Example 2.

24. A method as claimed in any one of claims 1 to 15, substantially as described in either of Example 1 or Example 2.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 93/01989

A. CLASSIFICATION OF SUBJECT MATTER

IPC 5 C12Q1/04 C12Q1/70 C12Q1/66

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JOURNAL OF CLINICAL MICROBIOLOGY vol. 20, no. 6, December 1984, WASHINGTON, D.C., US pages 1122 - 1125 SUSAN W. COOPER ET AL. 'Differentiation of Bacteroides ovatus and Bacteroides thetaiotamicon by Means of Bactriophage' see the whole document ---	1-24
A	JOURNAL OF FOOD PROTECTION vol. 47, no. 5, May 1984, AMES, IOWA, US pages 388 - 390 DWIGHT C. HIRSH AND LORI D. MARTIN 'Rapid Detection of Salmonella in Certified Raw Milk by Using Charge-Modified Filters and Felix-01 Bacteriophage' cited in the application see the whole document ---	1-24
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

22 December 1993

Date of mailing of the international search report

10. 01. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Doepfer, K-P

INTERNATIONAL SEARCH REPORT

Inter. Natl Application No

PCT/GB 93/01989

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>ONDERSTEEPOORT JOURNAL OF VETERINARY RESEARCH vol. 56 , 1989 , PRETORIA, SOUTH AFRICA pages 263 - 269 MARTHA L. VAN DER WALT AND HELENA C. STEYN 'The Biochemical Differentiation between Salmonella and Citrobacter' see the whole document ---</p>	1-24
A	<p>JOURNAL OF CLINICAL MICROBIOLOGY vol. 30, no. 3 , March 1992 , WASHINGTON, D.C., US pages 590 - 594 XIAOQING HE AND RUONAN PAN 'Bacteriophage Lytic Patterns for Identification of Salmonellae, Shigellae, Escherichia coli, Citrobacter freundii, and Enterobacter cloacae' see the whole document ---</p>	1-24
A	<p>PHYTOPATHOLOGY vol. 71, no. 3 , March 1981 , ST. PAUL, US pages 274 - 276 K.W. LIEW AND A.M. ALVAREZ 'Phage Typing and Lysotype Distribution of Xanthomonas campestris' see the whole document ---</p>	1-24
A	<p>GB,A,2 213 261 (THE PUBLIC HEALTH LABORATORY SERVICE BOARD) 9 August 1989 cited in the application see the whole document ---</p>	1-24
A	<p>WO,A,90 01559 (LONDON BIOTECHNOLOGY LIMITED) 22 February 1990 see the whole document & GB,A,2 240 845 (LONDON BIOTECHNOLOGY LTD.) cited in the application ---</p>	1-24
A	<p>US,A,4 348 478 (EIICHI NAKANO ET AL.) 7 September 1982 cited in the application -----</p>	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 93/01989

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A-2213261	09-08-89	AU-A- 2811789 EP-A- 0396584 WO-A- 8905356 JP-T- 3502521	05-07-89 14-11-90 15-06-89 13-06-91
WO-A-9001559	22-02-90	AU-B- 633507 AU-A- 4061289 EP-A- 0430987 GB-A- 2240845 JP-T- 4500005 WO-A- 8911544	04-02-93 05-03-90 12-06-91 14-08-91 09-01-92 30-11-89
GB-A-2240845	14-08-91	AU-B- 633507 AU-A- 4061289 EP-A- 0430987 WO-A- 9001559 JP-T- 4500005 WO-A- 8911544	04-02-93 05-03-90 12-06-91 22-02-90 09-01-92 30-11-89
US-A-4348478	07-09-82	JP-C- 1371400 JP-A- 55058096 JP-B- 61037917 GB-A,B 2042554	07-04-87 30-04-80 26-08-86 24-09-80